

GENERALIZED β -EXPANSIONS, SUBSTITUTION TILINGS, AND LOCAL FINITENESS

NATALIE PRIEBE FRANK AND E. ARTHUR ROBINSON, JR.

ABSTRACT. For a fairly general class of two-dimensional tiling substitutions, we prove that if the length expansion β is a Pisot number, then the tilings defined by the substitution must be locally finite. We also give a simple example of a two-dimensional substitution on rectangular tiles, with a non-Pisot length expansion β , such that no tiling admitted by the substitution is locally finite. The proofs of both results are effectively one-dimensional and involve the idea of a certain type of generalized β -transformation.

1. INTRODUCTION

This paper was motivated by the study of simple tiling substitutions such as the substitution S shown in Figure 1. To the left of the arrows, four prototiles




FIGURE 1. An inflate-and-subdivide rule S .

are shown along with their dimensions. The larger dimension, $\beta = (1 + \sqrt{13})/2 \approx 2.3027756377$, is a root of the polynomial $p(x) = x^2 - x - 3$. The patches to the right of the arrows show the application of the substitution S to each prototile. The substitution S can be applied repeatedly, to produce arbitrarily large patches of tiles. The first few applications, starting with the large square prototile, are shown in Figure 2. Our interest in S arises because its length expansion β is not a Pisot number. An algebraic integer β is called a *Pisot number* if $\beta > 1$, and all its Galois conjugates β' satisfy $|\beta'| < 1$. Here, β is not Pisot since its conjugate is $\beta' = (1 - \sqrt{13})/2 \approx -1.3027756377$.

It is known that whether or not the linear expansion of a tiling substitution S is a Pisot number has implications for the tiling dynamical system that it defines. (We make the distinction between the linear expansion and the length expansion of a substitution tiling precise in Section 4.) Solomyak proved in [8] that a one- or two-dimensional self-similar tiling dynamical system with real linear expansion β is weakly mixing if and only if β is not Pisot. A similar result holds for a two-dimensional self-similar tiling dynamical system with complex linear expansion $\lambda \in \mathbb{C} \setminus \mathbb{R}$. Solomyak showed that such a dynamical system is weakly mixing if and

¹2000 *Mathematics Subject Classification.* Primary 52C20; Secondary 37B50.

²*Key words and phrases.* Substitution sequence, self-similar tiling

only if λ is not a “complex Pisot” number. However, in the two-dimensional case the proofs of these results, as well as many similar results, use the additional assumption of “local finiteness” (a property that is automatic for the one-dimensional case). A tiling is defined to be *locally finite*¹ if, up to rigid motions, it contains only finitely many two-tile configurations.

For substitution tilings, local finiteness follows from “two-tile closure property” [5]. On the other hand, Danzer has conjectured [1] that non-locally finite tilings may in fact be generic for certain classes of substitutions. We will show that tilings admitted by the substitution in Figure 1 fail to be locally finite. It remains unclear the extent to which results like those in [8] can be extended to the non-locally finite case.

Although some examples of non-locally finite tilings have already appeared (see [1], see also [6]), our example has the benefit of simplicity. It belongs to a family [2] of tilings that can be created quite easily from any one-dimensional substitution. Moreover, our proof is based on a one-dimensional method that may be of independent interest.

To see how the lack of local finiteness appears from Figure 1, consider Figure 2, which shows three iterations of the large square tile. In the first iteration, we have circled a point where there is, in Danzer’s terminology [1], a “misfit situation”. Under the inflate-and-subdivide rule S the tiles are inflated by the linear map




FIGURE 2. Three iterations of the larger square tile.

βI ; we can follow the circled point’s location under this linear map. Following the circled point through two more iterations, we see that the misfit it creates moves around relative to the tile below it. We will prove that this point occupies infinitely many locations with respect to the tile below it. Therefore the infinite tiling generated by this process is not locally finite.

Now one can begin to see why the arguments are one-dimensional. Inspection of Figure 1 shows that there are two kinds of upper edges: a long one and a short one, which we denote l and s respectively. The reader can verify that a long upper edge

¹This property is also sometimes also called “finite local complexity”. In the theory Delone sets, however, “local finiteness” usually refers to a weaker property.

is substituted by a long followed by three short edges, ($l \rightarrow lsss$), and a short upper edge is substituted by a long one, ($s \rightarrow l$). The circled point in Figure 2 is always on the corner of two of the tiles above it, but it is (we will show) always in the interior of an upper edge. Thus we restrict our attention to the one-dimensional substitution on the upper edges to prove our result. In fact, following the misfit through these iterates is analogous to applying a suitably defined “generalized β -transformation”, which we discuss in the next section.

The organization of this paper is as follows. We first focus on the one-dimensional theory, which is itself of independent interest. In Section 2, we give the one-dimensional definition of a substitution tiling and a generalized β -transformation, and we show how they are related. In Section 3 we give conditions that guarantee finite or infinite orbits of the generalized β -transformations and relate this result to one-dimensional tilings. In Section 4 we apply this machinery to tilings of \mathbb{R}^2 .

2. DEFINITIONS

2.1. One-dimensional tiling substitutions. Let $\mathcal{A} = \{1, \dots, m\}$ and let \mathcal{A}^* denote the set of all (nonempty) finite words in \mathcal{A} . A *substitution* is mapping $\sigma : \mathcal{A} \rightarrow \mathcal{A}^*$. Given a substitution σ , the *structure matrix* is the $d \times d$ matrix M with entries $M_{i,j}$ equal to the number of occurrences of the symbol j in the word $\sigma(i)$. We require that M has a real eigenvalue $\beta > 1$ such that the right eigenvector $\vec{h} = (h_1, h_2, \dots, h_m)$ has strictly positive entries. It is worth noting that Perron-Frobenius theory ensures this in two special cases: irreducibility and primitivity. We say M is irreducible if for each i, j there is an n for which $(M^n)_{ij} > 0$. If a single n works for all i, j then we say M is primitive. By the Perron-Frobenius theorem, if M is irreducible then there exists an $r > 0$ for which $\vec{h} \in (r\mathbb{Q}(\beta))^m$.

Given a substitution σ , we construct a one-dimensional tiling substitution S and a corresponding tiling τ of the line. The *prototiles* $\mathcal{P} = \{P_1, \dots, P_m\}$ are intervals $P_j = [0, h_j]$ with lengths given by the entries of \vec{h} . A *tile* is a translation of a prototile $T = P_j + s = [s, s + h_j]$. A *tiling patch* π is a finite sequence of tiles laid end to end:

$$(1) \quad \pi = (P_{j_1} + s_1, P_{j_2} + s_2, \dots, P_{j_n} + s_n)$$

where

$$s_{i+1} = s_i + h_{j_i}.$$

Equivalently, in a tiling patch, the tiles are disjoint and the *support*, defined

$$(2) \quad \text{supp}(\pi) := \bigcup_{i=0}^n P_{j_i} + s_i,$$

is an interval. A *tiling* τ is an infinite set of disjoint tiles whose support is \mathbb{R} .

We call s the left endpoint of the tiling patch π , and if $\text{supp}(\pi) = [s, t]$, we call t the right endpoint. We can specify a tiling patch π by giving its left end point s and the word $\mathbf{w} = j_1 j_2 \dots j_n$ that specifies the prototiles that occur in its sequence. In particular, the patch in (1) is given by $\pi = \pi(s, \mathbf{w})$.

Now we can define the *tiling substitution* S corresponding to σ . It is the mapping from tilings to tiling patches defined by

$$S(P_j + s) = \pi(\beta s, \sigma(j)).$$

Clearly we have $\text{supp}(S(T)) = \beta \text{supp}(T)$, and $S(T + s) = S(T) + \beta s$. This allows us to extend S to a mapping on tiling patches, and even to tilings. A tiling τ is said to be S -admissible if every patch $\pi \subset \tau$ is equivalent to a patch of $\pi' \subset S^n P$ for

some $P \in \mathcal{P}$, $n \in \mathbb{N}$. The set of all S -admissible tilings (which is always nonempty) is called a *tiling space*, and is written X_S .

Example 1. Consider the substitution $l \rightarrow lsss$, $s \rightarrow l$ from the introduction. The structure matrix is

$$M = \begin{bmatrix} 1 & 3 \\ 1 & 0 \end{bmatrix},$$

which has characteristic polynomial $p(x) = x^2 - x - 3$, and the Perron eigenvalue $\beta = (1 + \sqrt{13})/2$. Taking $\vec{h} = (\beta, 1)$ as a positive right eigenvector, the prototiles are $P_1 = [0, \beta]$ and $P_2 = [0, 1]$. These tiles have the same lengths as the edges of the tiles in Figure 1. We will continue our discussion of this example below.

2.2. Generalized β -transformations. Fix an interval $E = [0, B)$ and a real number $\beta > 1$.

Definition 2.1. We say a mapping $F_\beta : E \rightarrow E$ is a generalized β -transformation if (i) F_β is piecewise linear, (ii) F_β is continuous from the right and (iii) $F'_\beta(x) = \beta$ for almost every $x \in I$.

More explicitly, F_β is a generalized β -transformation if there is a partition of E into l subintervals $I_j = [x_{j-1}, x_j)$, with endpoints $0 = x_0 < x_1 < \dots < x_l = B$, and real numbers y_1, y_2, \dots, y_l with $\beta I_j - y_j \subseteq E$ for $j = 1, \dots, l$, so that

$$(3) \quad F_\beta(x) = \beta x - y_j \text{ whenever } x \in I_j.$$

One example of this is the classical β -transformation $T_\beta : [0, 1) \rightarrow [0, 1)$, defined $T_\beta(x) = \beta x \bmod 1$ (see Renyi [4]). A type of generalized β -transformation, more general than what we consider here, was studied by Wilkinson [10]. He showed under several additional conditions that F_β has an absolutely continuous invariant measure and is weakly Bernoulli with respect to the partition into the intervals I_j .

Definition 2.2. We call a generalized β -transformation F_β algebraic if β is an algebraic number and there is an $r \in \mathbb{R}$ for which

$$x_0, \dots, x_l, y_1, \dots, y_l \in r\mathbb{Q}(\beta).$$

2.3. Substitution tilings and generalized β -transformations. Now let us think of the circled point in Figure 2 as being the origin situated in the interior of the one-dimensional tile that is the upper edge of a two-dimensional tile (Figure 3).




FIGURE 3. The map \bar{S} finds the central tile after the substitution is applied.

Suppose that $T = [s, t)$ is a tile with $s \leq 0$ and $t > 0$, so that $0 \in T$. We call such a tile a *central tile*, illustrated in Figure 3 with $T = [1 - \beta, 1)$. It follows that $0 \in \text{supp}(S(T)) = [\beta s, \beta t)$, and thus there is a unique central tile in $S(T) = (P_{j_1} + s_1, P_{j_2} + s_2, \dots, P_{j_n} + s_n)$, i.e., $0 \in P_{j_i} + s_i$ for some i . We can therefore

define a mapping \bar{S} from the set of all central tiles to itself by taking $\bar{S}(T)$ to be the central tile in $S(T)$. We are interested in the effects of iterating \bar{S} on central tiles, but it is convenient to think of this process a little differently.

First we consider the patch $\pi := \pi(0, 123\dots m) = (E_1, E_2, \dots, E_m)$ where $E_j = J_j + g_j$ with $g_0 = 0$ and $g_{j+1} = g_j + h_j$. We write the support of π as $E = [0, B]$. The mapping $F : E \rightarrow E$ is defined as follows. For $x \in E$, find $j \in \mathcal{A}$ so that $x \in E_j$. Then $E_j - x$ is a central tile, and $\bar{S}(E_j - x) = E_k - y$ for some unique $k \in \mathcal{A}$ and $y \in \mathbb{R}$. We put $y = F(x)$, noting that $F(x) \in E_k$.

We may write F explicitly as a generalized β -transformation by partitioning each E_j into subintervals that will be sent by F onto whole tiles (the canonical subdivision suggested by S). We obtain a partition $0 = x_0 < x_1 < \dots < x_l = B$ into intervals I_1, \dots, I_l such that for each I_k there is a j such that $F(I_k) = E_j$. The restriction of F to I_k is the linear map taking I_k onto E_j . So for $x \in I_k$ we have

$$(4) \quad F(x) = \beta(x - x_{k-1}) + g_{j-1};$$

hence $y_k = -\beta x_{k-1} + g_{j-1}$.

Example 2. The generalized β -transformation for Example 1 is defined on the interval $E = [0, 1 + \beta)$ so that $E_1 = [0, \beta)$ and $E_2 = [\beta, \beta + 1)$. The partition of E given by the substitution is $x_0 = 0, x_1 = 1, x_2 = 1 + 1/\beta, x_3 = 1 + 2/\beta, x_4 = 1 + 3/\beta = \beta$, and $x_5 = 1 + \beta$. Since $g_0 = 0$ and $g_1 = \beta$ our translations are $y_1 = 0, y_2 = 0, y_3 = 1, y_4 = 2$, and $y_5 = 3 + \beta$. The transformation is as shown in Figure 4.




FIGURE 4. The map $F : E \rightarrow E$.

Since each interval in a tiling substitution expands by a factor of β , we have the following lemma.

Lemma 2.3. *The mapping F given by a tiling substitution with expansion constant β is an algebraic generalized β -transformation whenever $\vec{h} \in (r\mathbb{Q}(\beta))^m$.*

This is certainly true whenever M is irreducible or primitive.

3. ONE-DIMENSIONAL RESULTS

Suppose $\beta > 1$ is a root of the irreducible monic polynomial $p(x) = g_0 + g_1x + g_2x^2 + \dots + x^d \in \mathbb{Z}[x]$, where $d > 1$. We can identify $\mathbb{Q}(\beta)$ with \mathbb{Q}^d via the basis $\{1, \beta, \beta^2, \dots, \beta^{d-1}\}$ over \mathbb{Q} . Multiplication by β in $\mathbb{Q}(\beta)$ corresponds to multiplication by A in \mathbb{Q}^d , where

$$(5) \quad A = \begin{bmatrix} 0 & 0 & \cdots & -g_0 \\ 1 & 0 & \cdots & -g_1 \\ 0 & 1 & \cdots & -g_2 \\ \vdots & & & \\ 0 & \cdots & 1 & -g_{d-1} \end{bmatrix}$$

is the companion matrix of $p(x)$.

To fully understand the action of A , we embed \mathbb{Q}^d in \mathbb{C}^d (although if the eigenvalues are all real, it suffices to embed in \mathbb{R}^d). We define a surjective linear map $\phi : \mathbb{C}^d \rightarrow \mathbb{C}$ by

$$\phi(\vec{w}) = w_0 + w_1\beta + w_2\beta^2 + \dots + w_{d-1}\beta^{d-1}.$$

The restriction of ϕ to \mathbb{Q}^d is the isomorphism between \mathbb{Q}^d and $\mathbb{Q}(\beta)$. Moreover, for any $\vec{w} \in \mathbb{C}^d$,

$$(6) \quad \phi(A\vec{w}) = \beta\phi(\vec{w}).$$

The Galois conjugates $\beta_2, \beta_3, \dots, \beta_d$ of β are eigenvalues of A with eigenvectors $\vec{v}_2, \vec{v}_3, \dots, \vec{v}_d$. By linearity $\phi(A\vec{v}_j) = \phi(\beta_j\vec{v}_j) = \beta_j\phi(\vec{v}_j)$, but by (6) we also have $\phi(A\vec{v}_j) = \beta\phi(\vec{v}_j)$, and so $\phi(\vec{v}_j) = 0$ for all $j = 2, 3, \dots, d$. Let V_0 be the subspace of \mathbb{C}^d spanned by $\vec{v}_1, \dots, \vec{v}_d$, and let V^+ be the (one-dimensional) eigenspace of β . Since ϕ is not the zero map, we can select the vector $\vec{v} \in V^+$ with $\phi(\vec{v}) = 1$ as a basis vector.

Consider fixed an algebraic generalized β -transformation $F : E \rightarrow E$. Recalling the notation of Subsection 2.2 we have the partition of E given by $0 = x_0 < x_1 < \dots < x_l = B$ with $x_j \in \mathbb{Q}(\beta)$; we note that by definition $\beta(x_j - x_{j-1}) \leq B$. We denote the j th subinterval as $I_j = [x_{j-1}, x_j)$, and we have $y_1, y_2, \dots, y_l \in \mathbb{Q}(\beta)$ so that $\beta I_j - y_j \subseteq E$.

Since $\phi(\vec{u}) = 0$ for any $\vec{u} \in V_0$, we know that $\phi(c\vec{v} + \vec{u}) \in \mathbb{R}$ if and only if $c \in \mathbb{R}$. Thus we can define

$$W_{[a,b]} = \{\vec{w} : \phi(\vec{w}) \in [a, b]\} = \{c\vec{v} + \vec{u} : \vec{u} \in V_0 \text{ and } c \in [a, b]\} \subseteq \mathbb{C}$$

to be the set of all points in \mathbb{C}^d that ϕ maps onto the real interval $[a, b]$.

Let W_E denote the subset of \mathbb{C}^d corresponding to E and denote the subsets of \mathbb{C}^d that correspond to the intervals I_j as $W_j = W_{I_j}$; the W_j s form a partition of W_E . There are unique vectors $\vec{w}_1, \vec{w}_2, \dots, \vec{w}_d$ in \mathbb{Q}^d for which $\phi(\vec{w}_i) = y_i$, and one can easily check that $AW_j - \vec{w}_j \subseteq W_E$. Thus we can define the map $f : W_E \rightarrow W_E$ as

$$(7) \quad f(\vec{w}) = A\vec{w} - \vec{w}_j \text{ whenever } \vec{w} \in W_j;$$

and we see that

$$(8) \quad \phi \circ f = F \circ \phi.$$

Because our interest lies with the mapping F , the theorem below is stated in terms of it, but the result is actually proved for f and then extended to F . The proof is mostly linear algebra and uses ideas similar to [1], [7] and [9].

Theorem 3.1. Let $\beta, \beta_2, \dots, \beta_d, E$, and F be defined as above.

- (1) If $|\beta_j| < 1$ for all $j = 2, 3, \dots, d$, then for any $x \in \mathbb{Q}(\beta) \cap E$ the orbit of x under F is finite and therefore eventually periodic.
- (2) If $|\beta_j| > 1$ for some $j \in \{2, 3, \dots, d\}$, then there exists an $x \in \mathbb{Q}(\beta) \cap E$ such that the orbit of x under F is infinite.

Proof. Since $\phi \circ f = F \circ \phi$ and the map ϕ is a bijection when restricted to \mathbb{Q}^d it suffices to prove the results for f .

Suppose $|\beta_j| < 1$ for all $j = 2, 3, \dots, d$. Let $\vec{w} \in \mathbb{Q}(\beta) \cap W_E$ and \vec{w}_i from (7) be written in terms of the eigenvectors of A as

$$\vec{w} = c\vec{v} + \sum_{i=2}^d c_i \vec{v}_i \quad \text{and} \quad \vec{w}_j = w_{j,1}\vec{v} + \sum_{i=2}^d w_{j,i} \vec{v}_i.$$

For $i = 2, \dots, d$ choose

$$r_i > \frac{|w_{j,i}|}{1 - |\beta_i|} \quad \text{for all } j \in 1, 2, \dots, d.$$

Define the *trapping region* \mathcal{R}_{trap} to be the set of all points in W_E such that $c_i \leq r_i$.

We claim that \mathcal{R}_{trap} is invariant under the action of f . Suppose $\vec{w} \in \mathcal{R}_{trap}$ with such a choice of values, and find $j \in 1, 2, \dots, l$ such that $\vec{w} \in W_j$. Then

$$(9) \quad f(\vec{w}) = A\vec{w} - \vec{w}_j = (\beta\vec{v} + \sum_{i=2}^d \beta_i c_i \vec{v}_i) - (w_{j,1}\vec{v} + \sum_{i=2}^d w_{j,i} \vec{v}_i).$$

Since the component in the direction of \vec{v} is unrelated to the definition of \mathcal{R}_{trap} we see that we must have $|\beta_i c_i - w_{j,i}| \leq r_i$ for $i = 2, 3, \dots, d$. This is true since $|w_{j,i}| < (1 - |\beta_i|)r_i$ and $|c_i| \leq r_i$. This proves the invariance.

Now note that there is some $q \in \mathbb{Z}$ for which \vec{w} and the \vec{w}_j s are in $\frac{1}{q}\mathbb{Z}^d$. Since A is an integer matrix it leaves $\frac{1}{q}\mathbb{Z}^d$ invariant and so $f\left(\frac{1}{q}\mathbb{Z}^d \cap W_E\right) \subseteq \frac{1}{q}\mathbb{Z}^d \cap W_E$. Choose the numbers r_i for the trapping region so that $\vec{w} \in \mathcal{R}_{trap}$; clearly $f\left(\frac{1}{q}\mathbb{Z}^d \cap \mathcal{R}_{trap}\right) \subseteq \frac{1}{q}\mathbb{Z}^d \cap \mathcal{R}_{trap}$. Since $\frac{1}{q}\mathbb{Z}^d \cap \mathcal{R}_{trap}$ is a finite set the orbit of \vec{w} is finite and thus eventually periodic.

To prove the second part of the theorem we must show there is an infinite *escape region* $\mathcal{R}_{esc} \subset W_E$ for which orbits beginning in \mathcal{R}_{esc} tend to infinity. Without loss of generality we may assume that $|\beta_2| > 1$. Choose r_2 so that

$$r_2 > \frac{|w_{j,2}|}{|\beta_2| - 1} \quad \text{for all } j = 1, 2, \dots, d.$$

We define

$$\mathcal{R}_{esc} = \{\vec{w} = c\vec{v} + \sum_{i=2}^d c_i \vec{v}_i \subset W_E \text{ such that } |c_2| > r_2\}.$$

To prove the result, we must show that for any $\vec{w} \in \mathcal{R}_{esc} \cap \frac{1}{q}\mathbb{Z}^d$, the component of $f(\vec{w})$ in the direction of \vec{v}_2 exceeds that of \vec{w} in modulus. Supposing again that $\vec{w} \in W_j$, by the computation in equation (9) this amounts to showing that $|\beta_2 c_2 - w_{j,2}| > |c_2|$. By choice of r_2 we have that $|w_{j,2}| < (|\beta_2| - 1)r_2 \leq (|\beta_2| - 1)|c_2|$ and the result follows. \square

Corollary 3.2. *Let S be a one-dimensional tiling substitution with expansion constant $\beta > 1$ and $\vec{h} \in (r\mathbb{Q}(\beta))^m$. Let T be a central tile. (i) Suppose β is a Pisot number. If the endpoints of T belong to $\mathbb{Q}(\beta)$, then there exist $K, L \in \mathbb{N}$ so that $\bar{S}^{NL}(T) = \bar{S}^K(T)$ for all $N > 0$. (ii) Suppose β is non-Pisot. Then there is a $q \in \mathbb{N}$ and finite $E \subseteq \frac{1}{q}\mathbb{Z}_\beta$ such that for any $s \in \frac{1}{q}\mathbb{Z}_\beta \setminus E$, if $T = [-s, t)$ is central, then $\bar{S}^K(T) \neq \bar{S}^L(T)$ for any $K > L \geq 1$.*

4. TILING RESULTS

In this section we study a class of self-similar substitution tilings in $\mathbb{R}^2 \cong \mathbb{C}$. Our first result is that if a substitution S has a Pisot length expansion, then the tilings it produces are locally finite. This solves a conjecture of Danzer [1] for the type of tilings that we consider. In this section we also prove that tilings generated by the substitution from Figure 1 are not locally finite.

We begin with some general definitions. A *tile* T is a closed topological disk $D \subseteq \mathbb{C}$, together (possibly) with a label which is an element of some fixed finite set \mathcal{A} . We say $D = \text{supp}(T)$ is the *support* of T . Two tiles are called *equivalent* if they carry the same label and have congruent supports.

We will fix a finite set \mathcal{P} of inequivalent tiles, called *prototiles*, and assume all tiles T are equivalent to tiles from \mathcal{P} . A *packing* is a collection tiles that intersect only on their boundaries, and the support of a packing is the union of the supports of its tiles². A *tiling patch* π is a finite packing with a connected support. We denote the set of patches by \mathcal{P}^* . A *tiling* τ is a packing with support \mathbb{C} .

Let $\lambda \in \mathbb{C}$ with $|\lambda| > 1$. Suppose $S : \mathcal{P} \rightarrow \mathcal{P}^*$ is such that for each $P \in \mathcal{P}$, the $\text{supp}(S(P))$ is congruent³ to $\lambda \cdot \text{supp}(P) := \lambda \cdot P$. We call S a *self-similar tiling substitution* (or simply a *substitution* when there is no danger of confusion). We call λ the *linear expansion* of S , equating multiplication by λ with a linear map on \mathbb{R}^2 ; volumes must expand by a factor of $\lambda\bar{\lambda}$. We call $\beta = |\lambda|$ the *length expansion* of S because it gives the expansion of one-dimensional objects.

The substitution S extends in an obvious way to any tile T , and can thus be iterated to obtain infinitely large patches $S^n(P)$. Following the one-dimensional case, we say a tiling τ is *S -admissible* if for every patch $\pi \in \tau$ is equivalent to a patch $\pi' \in S^n(P)$ for some $P \in \mathcal{P}$ and $n \geq 0$. We denote the set of S -admissible tilings τ by X_S .

The following Lemma may be well-known, but we include a short proof for completeness.

Lemma 4.1. *Let \mathcal{P} be a finite prototile set and let S be a tiling substitution on \mathcal{P} . No prototile in \mathcal{P} contains a nontrivial circular arc in its boundary.*

Proof. Suppose the boundary of a prototile $P \in \mathcal{P}$ contains an arc of radius $r > 0$ subtending an angle $\theta > 0$. The boundary of $S^n(P)$ contains arcs of radius $\beta^n r$ subtending the angle θ , and this implies that there are prototiles in \mathcal{P} that have circular arcs of radius $\beta^n r$ in their boundaries. Since the prototile set is finite, we can find the maximum angle $\theta_n > 0$ subtended by the arc of radius $\beta^n r$ in any

²In the one dimensional case, it was convenient to take tiles as half-open and insist on disjointness. This also forced equivalence to preserve orientation, a restriction we do not impose here.

³In the event that there is more than one congruence between the supports $S(P)$ and $\lambda \cdot P$, we will use the implicit parameterization of the boundaries to ensure uniqueness of $S(T)$.

prototile. The arclengths $\theta_n \beta^n r$ must go to zero in order for the prototiles to be topological disks.

Let d_{max} and v_{min} be the maximum diameter and minimum volume of prototiles in \mathcal{P} , respectively. The tiles inside of $S^n(P)$ adjacent to the circular arc of length $\theta \beta^n r$ lie inside a strip of volume

$$\theta/2[(\beta^n r)^2 - (\beta^n r - d_{max})^2] + 2\pi d_{max}^2,$$

where $2\pi d_{max}^2$ ensures the ends of the strip are covered. The maximum number of tiles that can fit inside this strip is

$$\frac{\theta/2[2d_{max}\beta^n r - d_{max}^2] + 2\pi d_{max}^2}{v_{min}} < \frac{\theta d_{max}\beta^n r}{v_{min}} + K,$$

where $K = (2\pi - \theta/2)d_{max}^2/v_{min}$ does not depend on n . The minimum number of tiles required to cover the arc of length $\theta \beta^n r$ is $(\theta \beta^n r)/(\theta_n \beta^n r) = \theta/\theta_n$. For all $n = 1, 2, \dots$ we must have

$$\begin{aligned} \frac{\theta}{\theta_n} &\leq \frac{\theta d_{max}\beta^n r}{v_{min}} + K, \text{ or} \\ \theta &\leq \frac{\theta d_{max}}{v_{min}}(\theta_n \beta^n r) + K\theta_n. \end{aligned}$$

We have noted that both terms on the right must go to zero, forcing θ to be zero also. This contradiction completes the proof. \square

4.1. Local finiteness when the length expansion is Pisot. For the result we are going to prove, we will need some mild additional geometric hypotheses on prototiles and tilings. A *straight edge* ℓ in the boundary of a tile T is a line segment that cannot be extended without leaving the boundary. Any such edge has a left and a right *endpoint*, relative to the boundary being traversed clockwise. The absence of circular arcs explains the terminology in the following definition.

Definition 4.2. *A tile T is called fractagonal if:*

- (1) *T has only finitely many straight edges, and*
- (2) *for any straight edge ℓ of T , the rest of T lies in the interior of a half-plane generated by ℓ (we call this weak convexity).*

The authors conjecture that the first property is automatically satisfied by the prototiles of a self-similar tiling substitution.

We also need an assumption that for straight edges, collinearity implies comeasurability in the following sense:

Definition 4.3. *We say a tiling substitution S is a property-(C) substitution if whenever two straight edges ℓ_1 and ℓ_2 are collinear in a tiling $\tau \in X_S$, there is an $r > 0$ for which $|\ell_1|, |\ell_2| \in r\mathbb{Q}(\beta)$, where $\beta = |\lambda|$ is the length expansion of S .*

Theorem 4.4. *Suppose that S is a property-(C) tiling substitution on a finite set \mathcal{P} of fractagonal prototiles. If the length expansion of S is a Pisot number, then any tiling in $\tau \in X_S$ is locally finite.*

Notes. There are two common assumptions that are not necessary. The substitution need not be primitive in that the structure matrix M may never satisfy $M^n > 0$. Tilings in X_S need not be translationally finite, i.e. prototiles may appear in an infinite number of orientations. The authors are not aware of any primitive tiling substitutions that are not property-(C).

Proof. A *simple adjacency* in $\tau \in X_S$ is a two-tile patch $\alpha \subset \tau$ where the intersection is along straight edges. Kenyon [3] proves that for any tiling τ made from a finite prototile set \mathcal{P} , if τ has an infinite number of inequivalent two-tile patches, then those patches are either simple adjacencies, or they occur along an entire circle of tile boundaries. By Lemma 4.1 the latter cannot occur in a substitution tiling. It follows that in order to show $\tau \in X_S$ is locally finite, it suffices to show that there are only finitely many simple adjacencies α .

Fix a simple adjacency α in $\tau \in X_S$. Two straight edges define α ; the difference between two vertices, one selected from each edge, is called an *offset* of α . Thus each α defines four different offsets. It suffices to show that the total number of offsets is finite.

Let the offset \mathcal{O} of α be fixed. We want to see \mathcal{O} as the difference of sums of edge lengths of tiles (see Figure 6). As such, it is convenient to consider edges as being intervals on the positive real axis. We call a tile T placed below (resp. above) the real axis with a straight edge on the real axis a *lower* (resp. *upper*) tile. By applying a rigid motion, we can take edge representatives with their left (resp. right) endpoints at the origin. We call these *edge sets* \mathcal{E}_ℓ and \mathcal{E}_u .

By weak convexity, the substitution S induces a one-dimensional substitution S_ℓ on \mathcal{E}_ℓ in the sense of Subsection 2.1. It is clear that the length expansion β of S is the expansion of S_ℓ , so the one-dimensional substitution has a Pisot expansion constant. If T is the lower tile containing $e \in \mathcal{E}_\ell$, we simply rotate $S(T)$ so that $\lambda \cdot e$ is mapped to $\beta \cdot e$ and look at the lower tiles of $\beta \cdot e$. Similarly we can create the one-dimensional substitution S_u on \mathcal{E}_u by considering the substitution on upper tiles. In particular, S_u is just the reverse of S_ℓ .

By the construction of τ , there is some smallest $n \geq 0$ for which there exists a tile T such that α is a sub-patch of $S^{n+1}(T)$. The intersection of the two tiles in α is contained in a line segment L that is composed of tile edges and is maximal in the sense that it cannot be extended without either entering the interior of some tile in $S^{n+1}(T)$ or leaving $S^{n+1}(T)$ entirely. We call L a *fault line* through the tile $S^{n+1}(T)$. Since \mathcal{O} depends only on α , we can use L to compute the value of \mathcal{O} , even if the tiles surrounding L in $S^{n+1}(T)$ do not appear in τ (except of course those in α).

If $n = 0$, then $S(T)$ can be moved to the positive x -axis so that an endpoint of L lies at the origin. By the weak convexity assumption it follows that L can be seen as a union of edges of upper tiles and also as a union of edges of lower tiles. In this case L , along with the information about which level-one tile it is in and which tile edges define it, is called an *initial segment*.

If $n > 0$, then by the minimality of n , there must be two level- n tiles whose boundaries intersect α . From this we see that L must contain a line segment L' that is a union of edges of level- n tiles. Since the tiles are assumed to be fractagonal, weak convexity implies that if $S^{n+1}(T)$ is moved so that L' is on the x -axis, then L' is the union of edges of upper level- n tiles and also the union of edges of lower level- n tiles. Thus it is the image under S^n of some initial segment L'' .

Since there are only a finite number of level-one tiles up to equivalence, there are only a finite number of initial segments up to equivalence. Let \mathcal{L} be a set of representatives of these equivalence classes, and without loss of generality choose these representatives to be on the positive real axis with an endpoint at the origin. For each $L \in \mathcal{L}$, the endpoints of the lower tiles of L create a partition $0 = a_0 <$

$a_1 < \dots < a_{k_\ell} = |L|$, where k_ℓ is the number of lower tiles of L and $|L|$ is the length of L . Any such a_i for any initial segment L will be called a *prefix* of L . Similarly, the endpoints of the upper tiles of L create a partition $0 = b_0 < b_1 < \dots < b_{k_u} = |L|$ where k_u is the number of upper tiles of L . A *suffix* of an initial segment L is $|L| - b_i$, where b_i is in the partition of L . In the same manner, we may compute prefixes of $S_\ell(e)$ for $e \in \mathcal{E}_\ell$ and suffixes of $S_u(e)$ for $e \in \mathcal{E}_u$. Since $|S_\ell(e)| = |S_u(e)| = \beta|e|$, the prefixes will be partition elements of $[0, \beta|e|]$, and the suffixes will be $\beta|e|$ minus partition elements of $[0, \beta|e|]$. We collect all the possible prefixes into a prefix set \mathcal{Q}_{px} and all possible suffixes into a suffix set \mathcal{Q}_{sx} . It is important to note that both \mathcal{Q}_{px} and \mathcal{Q}_{sx} are finite.

Since S is a property-(C) substitution, without loss of generality we can assume that all straight edge lengths belong to $\mathbb{Q}(\beta)$. Since each $P \in \mathcal{P}$ is fractagonal, there are only a finite number of straight edges. Thus there is a $q \in \mathbb{N}$ so that all straight edge lengths are in $\frac{1}{q}\mathbb{Z}[\beta]$. This implies that the lengths of all initial segments, and the lengths of all prefixes and suffixes in \mathcal{Q}_{px} and \mathcal{Q}_{sx} are also in $\frac{1}{q}\mathbb{Z}[\beta]$. Since offsets are the difference between a sum of upper edges and a sum of lower edges, $\mathcal{O} \subset \frac{1}{q}\mathbb{Z}[\beta]$.

We have already seen that \mathcal{O} lies in a segment L' that is the image $\lambda^n L''$ for some initial segment L'' . Let L_0 be the representative in \mathcal{L} of L'' . The tiles in α , which intersect $L' = \lambda^n \cdot L''$, are equivalent to lower and upper tiles T_ℓ and T_u intersecting $\beta^n \cdot L_0$. In fact the edges of T_ℓ and T_u are lower and upper edges that are edges in $S_\ell(L_0)$ and $S_u(L_0)$ when we consider L_0 in terms of its lower and upper tiles. The endpoints of edges in α used to compute \mathcal{O} correspond to points $a \in T_\ell$ and $b \in T_u$ (see Figure 5), so to compute \mathcal{O} we need to compute $|b - a|$. Without loss of generality assume $a \leq b$.




FIGURE 5. The simple adjacency α is equivalent to $T_u \cup T_\ell$.

We may compute the left endpoint a in terms of powers of β applied to elements of \mathcal{Q}_{px} ; we do the same for the upper endpoint b using powers of β applied to elements of \mathcal{Q}_{sx} . (See Figure 6). Let $p_n \in \mathcal{Q}_{px}$ be the largest prefix of L_0 for which $\beta^n p_n \leq a$. If equality holds, we have computed a , otherwise we continue. We denote by e_{n-1} the lower edge immediately following p_n in L_0 . This edge will, after application of S_l^n , contain a (otherwise p_n wasn't maximal). So there is a largest prefix of e_{n-1} which we call $p_{n-1} \in \mathcal{Q}_{px}$ for which $\beta^n p_n + \beta^{n-1} p_{n-1} \leq a$. If equality holds, we are done and the remainder of the prefixes will be the zero prefix. If not, we have that p_{n-1} is a prefix of $S_\ell(e_{n-1})$ for some edge $e_{n-1} \in \mathcal{E}_\ell$, so the lower edge immediately following p_{n-1} in $S_\ell(e_{n-1})$ will, after application of S_l^{n-1} , contain a . We can continue in this fashion, choosing prefixes p_{n-2}, \dots, p_0 until we obtain that $a = \sum_{k=0}^n \beta^k p_k$.

Similarly we can compute the value of b . Let $s_n \in \mathcal{Q}_{sx}$ be the largest suffix of L_0 for which $b \leq \beta^n (|L_0| - s_n)$. If equality holds we are done and all subsequent suffixes are the zero suffix. If not we select the edge e_{n-1} immediately preceding

$|L_0| - s_n$ and find its suffix s_{n-1} so that $b \leq \beta^n(|L_0| - s_n) - \beta^{n-1}s_{n-1}$. We continue choosing suffixes s_{n-2}, \dots, s_0 until we obtain that $b = \beta^n|L_0| - \sum_{k=0}^n \beta^k s_k$.




FIGURE 6. The computation of a and b as seen in $[0, \beta^n |L_0|]$.

Thus the offset distance \mathcal{O} is given by

$$(10) \quad \mathcal{O} = \beta^n |L_0| + \sum_{k=0}^n \beta^k (-s_k - p_k).$$

Our interest, then, is in all possible combinations of suffixes and prefixes. We collect these into a set \mathcal{D} which we must show is finite.

If β is an integer, then the result follows immediately since \mathcal{O} is a bounded by the length of the longest straight edge and is an element of $\frac{1}{q}\mathbb{Z}$.

If β is not an integer, we need to use the ideas and notation of Section 3. Recall that the map $\phi : \mathbb{C}^d \rightarrow \mathbb{C}$ is a bijection between \mathbb{Q}^d and $\mathbb{Q}(\beta)$, and that multiplication by the matrix A in \mathbb{Q}^d is equivalent to multiplication by β in $\mathbb{Q}(\beta)$. We can find preimages $\vec{w}_k \in \mathbb{Q}^d$ such that $\phi(\vec{w}_k) = -s_k - p_k$ for $k = 0, \dots, n$. Since elements of \mathcal{D} are in $\frac{1}{q}\mathbb{Z}[\beta]$, we have that each \vec{w}_k must be in $\frac{1}{q}\mathbb{Z}^d$. Equation (10) becomes

$$(11) \quad \mathcal{O} = \beta^n |L_0| + \sum_{k=0}^n \beta^k \phi(\vec{w}_k) = \beta^n |L_0| + \phi\left(\sum_{k=0}^n A^k \vec{w}_k\right).$$

We write the preimages of \mathcal{D} in terms of the eigenvectors $\vec{v}_1, \dots, \vec{v}_d$. Since ϕ projects onto \vec{v}_1 sending all other eigenvectors to 0, and since \mathcal{O} is necessarily less than the length of the longest edge, there is an $m_1 > 0$ that is an upper bound on the modulus of the component of \mathcal{O} in the \vec{v}_1 direction. For any $j = 2, \dots, d$, denote by m_j the maximum modulus of the component in the \vec{v}_j direction of any element of the difference set \mathcal{D} . The modulus of the component of $\sum_{k=0}^n A^k \vec{w}_k$ in the \vec{v}_j direction is less than or equal to

$$\sum_{k=0}^n \beta_j^k m_j < \sum_{k=0}^{\infty} \beta_j^k m_j = \frac{m_j}{1 - \beta_j},$$

since $|\beta_j| < 1$ by assumption when $j = 2, \dots, d$. So we see that the vectors which project onto \mathcal{O} must lie in a bounded region. There are only a finite number of elements of $\frac{1}{q}\mathbb{Z}^d$ in a bounded region, so there are only a finite number of offsets \mathcal{O} , and hence only a finite number of simple adjacencies A . The tiling x is locally finite. \square

Comment. A complex algebraic integer λ is called a *complex Pisot number* if $|\lambda| > 1$ and all algebraic conjugates $\lambda' \neq \bar{\lambda}$ satisfy $|\lambda'| < 1$ (see [8], [5]). If λ is complex Pisot, $\beta = |\lambda|$ may not be real Pisot (see [5] for an example). We do not know whether Theorem 4.4 always holds in the case of a complex Pisot linear expansion λ .

4.2. The Figure 1 example is not locally finite. Let a denote the large square prototile from Figure 1. Since the substitution of any tile includes a tile of type a , it follows that a block of type $S^n(a)$ must appear in $S^{n+1}(e)$ for any tile type and for any $n = 1, 2, \dots$ and so every tiling $x \in X$ must contain blocks of type $S^n(a)$ for all $n = 1, 2, \dots$

The circled point in Figure 2 appears inside the block $S(a)$. This point represents a vertex of the tiles above it and it intersects the upper boundary of the tile below it at a point $\beta - 1$ units from the left endpoint of that boundary. Each time we apply the substitution, the circled point determines the offset between the tile below it and the two tiles above it. This offset is completely determined by F . If we can show that the orbit of $\beta - 1$ is infinite then this implies that there are an infinite number of offsets and hence the tiling is not locally finite. It is convenient to compute this orbit for f rather than F .

Example 2 gives the algebraic generalized β -transformation associated with the substitution on upper edges of tiles. Noting that the eigenvalues are real, we compute the corresponding action f in $W_E \subset \mathbb{R}^2$. The companion matrix of β is

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 3 \end{bmatrix}.$$

We choose as the generator of V^+ the vector $\vec{v} = (-\beta_2, 1)$ and as generator of V_0 we take $\vec{v}_2 = (-\beta, 1)$. The vectors mapping onto y_1, y_2, y_3, y_4 and y_5 are given by $\vec{w}_1 = (0, 0), \vec{w}_2 = (0, 0), \vec{w}_3 = (1, 0), \vec{w}_4 = (2, 0)$, and $\vec{w}_5 = (3, 1)$. The sets W_j have boundaries given by $V_0 + \phi^{-1}(x_{j-1})$ and $V_0 + \phi^{-1}(x_j)$, where the inverse is taken in \mathbb{Q}^2 . It is not difficult to check that the appropriate translation vectors are $(0, 0), (1, 0), (2/3, 1/3), (1/3, 2/3), (0, 1)$ and $(1, 1)$. (In this example one can also find these by looking at $A^{-1}\vec{w}_j$.) The regions W_j are as shown in Figure 7, along with the main eigenspace V^+ .




FIGURE 7. The regions W_j in the domain W_E of f .

We now establish an f -orbit which enters the escape region and thus tends to infinity. By our theorem, this orbit corresponds to an infinite F -orbit. We begin by computing the escape region. Looking in the proof of Theorem 3.1 (2), we see

that we must choose r_2 so that

$$r_2 > \frac{|w_{j,2}|}{|\beta_2| - 1} \quad \text{for all } j = 1, 2, 3, 4.$$

We recall that $\vec{w}_1 = (0, 0)$, $\vec{w}_2 = (1, 0)$, $\vec{w}_3 = (2, 0)$, and $\vec{w}_4 = (3, 1)$ and compute their components in the V_0 direction:

$$w_{1,2} = 0, \quad w_{2,2} = \frac{-1}{\sqrt{13}}, \quad w_{3,2} = \frac{-2}{\sqrt{13}}, \quad w_{4,2} = \frac{-7 + \sqrt{13}}{2\sqrt{13}}.$$

The largest of these in modulus is $|w_{3,2}| \approx .5547$. We see that

$$r_2 > \frac{|w_{3,2}|}{|\beta_2| - 1} \approx 1.8321,$$

so we simply choose $r_2 = 2$ to define \mathcal{R}_{esc} .

We will show that $\vec{w} = \phi^{-1}(\beta - 1) = (-1, 1)$ enters the escape region by computing the first several elements of its orbit. (This was made easier by using a cobwebbing program in Matlab to determine in which W_j the point was at each step.) Now $(-1, 1) \in W_1$, so $f(-1, 1) = A(-1, 1) - \vec{w}_1 = (3, 0)$. One can compute that $(3, 0) \in W_4$, so $f(3, 0) = A(3, 0) - \vec{w}_4 = (-3, 2)$. Continuing in this fashion the next several iterates are $(5, -1), (-6, 3), (9, -3), (-11, 6), (15, -6)$. The orbit enters the escape region at $(-6, 3)$ and the iterates begin to grow quite quickly. It is instructive and surprising to draw the region W_E , which is infinitely long and quite thin, and plot these points which must remain inside but which bounce further and further away from the origin.

REFERENCES

1. L. Danzer, Inflation species of planar tilings which are not of locally finite complexity *Proc. Steklov Inst. Math.* **239** (2002), pp. 118-126.
2. N. P. Frank, Non constant length \mathbb{Z}^d substitutions, in preparation.
3. R. Kenyon, Rigidity of planar tilings, *Invent. math.* **107** (1992), 637-651.
4. A. Renyi, Representations for real numbers and their ergodic properties, *Acta Math. Acad. Sci. Hungar* **8** (1957), 477-493.
5. E. A. Robinson, Symbolic dynamics and tilings of \mathbb{R}^d , *Proceedings of Symposia in Applied Mathematics* **60** (2004), 81-119.
6. L. Sadun, Some generalizations of the Pinwheel Tiling, *Discrete Comp. Geom.* **20**, no.1 (1998), 79-110.
7. K. Schmidt, On periodic expansions of Pisot numbers and Salem numbers, *Bull. London Math. Soc.* **12** (1980), 269-278.
8. B. Solomyak, Dynamics of Self-Similar Tilings, *Ergodic Theory Dynamical Systems* **17** (1997), 695-738. Errata, *Ergodic Theory Dynamical Systems* **19** (1999), 1685.
9. W. Thurston, *Groups, Tilings, and Finite State Automata*, AMS Colloquium Lecture Notes, American Mathematical Society, Boulder, 1989.
10. K. M. Wilkinson, Ergodic properties of a class of piecewise linear transformations *Z. Wahr. verw. Gebiete* **31** (1975), pp. 303-328.

NATALIE PRIEBE FRANK, DEPARTMENT OF MATHEMATICS, VASSAR COLLEGE, BOX 248, POUGHKEEPSIE, NY 12604, NAFRANK@VASSAR.EDU

E. ARTHUR ROBINSON, JR., DEPARTMENT OF MATHEMATICS, GEORGE WASHINGTON UNIVERSITY, WASHINGTON, DC 20052, ROBINSON@GWU.EDU